



| | | |
|---|--|--|
| ОҢТҮСТІК-ҚАЗАҚСТАН MEDISINA AKADEMIASY «Оңтүстік Қазақстан медицина академиясы» АҚ |  | SOUTH KAZAKHSTAN MEDICAL ACADEMY АО «Южно-Казахстанская медицинская академия» |
| Department of «Pediatrics-1» | 044-38 | |
| Methodological guidelines | 1 page of 8 | |

**Methodological guidelines for students on teaching clinical skills in The center of practical skills
(using action algorithms)**

Specialty: «General medicine» 6B10101
Discipline: «Basics of childhood diseases-1»
Course: 4th course
Department: Pediatrics-1
Compiler: : Baltabaeva B.S.


Shymkent 2022 year

| | | |
|---|---|--|
| OÑTÚSTIK-QAZAQSTAN MEDISINA AKADEMIASY «Оңтүстік Қазақстан медицина академиясы» АҚ |  SKMA -1979- | SOUTH KAZAKHSTAN MEDICAL ACADEMY АО «Южно-Казахстанская медицинская академия» |
| Department of «Pediatrics-1» | 044-38 | |
| Methodological guidelines | 2 page of 8 | |

Considered at a meeting of the Department of Pediatrics-1

Protocol № 6 from « 24 »01 2023 year

Head of department, PhD  _____ Kemelbekov K.S.

| | | |
|---|--|--|
| ОҢТҮСТІК-ҚАЗАҚСТАН MEDISINA АКАДЕМИАСЫ «Оңтүстік Қазақстан медицина академиясы» АҚ |  | SOUTH KAZAKHSTAN MEDICAL ACADEMY АО «Южно-Казахстанская медицинская академия» |
| Department of «Pediatrics-1» | 044-38 | |
| Methodological guidelines | 3 page of 8 | |

1. The purpose of learning:

- correlation of symptoms and examination data with a care of a child with convulsive syndrome;
- determining a sequence of interventions in assisting a child with convulsive syndrome.

2. Theoretical knowledge necessary for mastering skills:

Convulsions are sudden attacks of clonic or clonic-tonic involuntary muscle contractions with or without loss of consciousness.

Classification:

- benign convulsions;
- epileptic;
- convulsions in infectious diseases;
- convulsions due to metabolic disorders (hypoglycemia, hyponatremia);
- convulsions caused by noncommunicable diseases (trauma, poisoning, diabetic ketoacidosis, glomerulonephritic encephalopathy).

Risk factors:

- burdened obstetric history (complicated birth, birth injury);
- hypoglycemia;
- severe fever;
- epilepsy in a personal and family anamnesis;
- infectious diseases of the Central nervous system (meningitis, encephalitis);
- head injury;
- poisoning with drugs or toxic drugs;
- post-vaccination complications;
- low levels of calcium, pyridoxine in the blood.

Diagnostic criterions:

- involuntary muscle contraction;
- tense or bulging fontanel;
- unnatural posture;
- signs of a head injury or other injury.


The list of the main diagnostic measures:

1. General blood test (3 parameters);
2. General urine test.

Tactics of treatment:

It is necessary to correctly position a child:

- If there is no suspicion of a neck injury:
 - turn the child on his side to reduce the risk of aspiration;
 - keep the child's head slightly turned and stabilize it by putting the cheek on the hand;
 - bend one leg of the child to stabilize the position of the body.
- If there is suspicion of a neck injury:

| | | |
|---|---|--|
| ОҢТҮСТІК-ҚАЗАҚСТАН MEDISINA АКАДЕМИАСЫ «Оңтүстік Қазақстан медицина академиясы» АҚ |  | SOUTH KAZAKHSTAN MEDICAL ACADEMY АО «Южно-Казахстанская медицинская академия» |
| Department of «Pediatrics-1» | 044-38 | |
| Methodological guidelines | 4 page of 8 | |

- stabilize the child's neck and leave him lying on his back;
- fix the child's forehead to a hard surface to secure the position;
- place bags of warm liquid on both sides of the baby's head to prevent head turns.
- when vomiting, turn the baby on its side, keeping the head in line with the torso.

In the presence of fever (38,5 and more) - paracetamol * 10-15 mg / kg and / or wipe the child with water at room temperature, do not prescribe oral drugs before relief of seizures due to the danger of aspiration.

Anticonvulsant treatment:

- diazepam * 0.5 mg / kg rectally or intramuscularly in infants < 2 months or intravenously diazepam 0.2-0.3 mg / kg If there is no effect – after 10 min. after administration – repeat. The maximum dose is 10 mg (with a body weight of 20 kg). For children < 2 months. - phenobarbital solution 20 mg / kg. Repeat after 30 min., if the cramps continue. Urgently to send to hospital.

List of essential medicines:

1. *paracetamol 500 mg, tablets;
2. *diazepam 5 mg, tablets; 10 mg/2ml, ampoules.

List of additional medicines:

1. phenobarbital 20 mg/5 ml, ampoules.

Indications for hospitalization:

A presence of seizures is an absolute indication for hospitalization.

Indicators of treatment effectiveness: relief of convulsive syndrome.

3. Essential clinical skills:


- counting respiratory, heart rate;
- auscultation of lungs, heart.

4. Clinical case:

A child is brought to a hospital in an ambulance. The child's mother came home from work on the call of a nanny. The nanny told her that the child vomited 3 times and he was very sleepy. The child's mother called an ambulance when the child lost consciousness in arms of the nurse and stopped breathing. The child a few days before kept a slight temperature, and had a runny nose. According to an ambulance crew report, the child had apnea when they arrived, the mother trying mouth-to-mouth CPR. By this time the child did not react to anything, he had cyanosis and bradycardia. Artificial ventilation and oxygen supply were started. The ambulance could not establish intravenous access and they took the child to the hospital.

The child begins to convulse.

Admission to the emergency room the baby has the following vital signs: heart rate is about 180, pressure 50/40, respiratory rate-40, SpO2 is not determined and temperature 38C. Fine bubbling rales are heard, hypoactive sounds of peristalsis. Heart rate - sinus tachycardia. A small bruise with a diameter of 2 cm. Abundant purulent discharge from a nose. A skin is covered with spots, a filling speed of capillaries is 5

| | | |
|---|--|--|
| ОҢТҮСТІК-ҚАЗАҚСТАН MEDISINA AKADEMIASY «Оңтүстік Қазақстан медицина академиясы» АҚ |  | SOUTH KAZAKHSTAN MEDICAL ACADEMY АО «Южно-Казакстанская медицинская академия» |
| Department of «Pediatrics-1» | 044-38 | |
| Methodological guidelines | 5 page of 8 | |

seconds. A frontal fontanel is swollen, pupils react to stimulus. Although all extremities move, the child is in a lethargic state.

Test results:

Complete blood count: hemoglobin 9.1, hematocrit 28.2%, white blood cells 14, platelets 124,000.

Biochemical blood test: Na 138, K 3.6, Cl 105, BUN 18, creatinine 0.5, glucose 74.

Respiratory syncytial virus test: negative.

Chest X-ray: clean.


5. Tasks:

In this scenario, 3 stages, a transition from one stage to another is carried out manually at a discretion of a mentor. In the first stage, a student examines a child in an emergency room and begins treatment. The child begins to convulse. In the final stage, students must examine the child for abuse and transfer the child to the children's intensive care department. When students conduct correct and timely interventions, the mentor should move on to the next stage.

In Stage 1 Admission to the emergency room, the baby has the following vital signs: heart rate is about 180, pressure is 50/40, respiratory rate is 40, SpO₂ is not determined and temperature is 38C. Fine bubbling rales are heard, hypoactive sounds of peristalsis. Heart rate - sinus tachycardia. Eyes are closed, a small bruise with a diameter of 2 cm is above the right eye. Abundant purulent discharge from the nose. A skin is covered with spots, a filling speed of capillaries is 5 seconds. A frontal fontanel is swollen, pupils react to a stimulus. Although all extremities move, the child is in a lethargic state. It is expected that a cadet will conduct a complete initial examination of the child, analyze the data, install a heart monitor and request an interpretation of the data. Also, the cadet must determine the correct dose and start infusion therapy (20 ml / kg = 72 ml), constantly monitor the infusion pump and the condition of the child, monitor the fluid flow rate and request the necessary laboratory tests. The cadet must also correctly and effectively communicate with the patient and relatives, satisfying their psychosocial needs and supporting them. The cadet must request the results of laboratory tests and X-ray images, analyze the data. Mentor must report test results.

In Stage 2 Convulsions 30 minutes later, the child's condition worsens, the baby's heart rate is about 200, the pressure is 80-90/50, the respiratory rate is 10-15, SpO₂ 80 with oxygen 2 l/m, the baby is observed for 1 minute tonic-clonic convulsions. Near-mouth cyanosis, pupils displaced to the left. It is expected that the cadet will ensure the safety of the child and the environment, control seizures and fill out the necessary documents, increase the oxygen supply to increase saturation, immediately inform the attending physician of an attack of seizures and receive new instructions. The cadet must re-examine, analyze the results and fill out the documents.

In Stage 3 Examination for abuse, the child's condition improves, heart rate 100, blood pressure 70/40, respiratory rate 30, SpO₂ 90 on oxygen 4 l/m. The heart rhythm is normal sinus, bilateral fine-bubbly moist rales are heard in lungs, and the child is in a lethargic

| | | |
|---|--|--|
| ОҢТҮСТІК-ҚАЗАҚСТАН MEDISINA AKADEMIASY «Оңтүстік Қазақстан медицина академиясы» АҚ |  | SOUTH KAZAKHSTAN MEDICAL ACADEMY АО «Южно-Казахстанская медицинская академия» |
| Department of «Pediatrics-1» | 044-38 | |
| Methodological guidelines | 6 page of 8 | |

state. It is expected that the cadet will correctly calculate the dose and inject intravenously phenytoin, according to 6 rules (15 mg x 3.6 kg = 54 mg = 1.08 ml for 5-15 minutes), establish monitoring of blood pressure, heart rate and pulse during the infusion of phenytoin, give parents to sign an agreement to conduct a lumbar puncture, assist in carrying out a lumbar puncture, raise the headboard by 30 degrees, make sure that the head position is neutral, limit the stimulation of the child to a minimum (silence, dimmed light).

6. Literature:

1. Лечение судорожных состояний и эпилепсии у детей. КПП. 2004 г. <http://medi.ru>.
2. Itimier, L. (2008). Shaken baby syndrome. The Journal or Perinatal & Neonatal Nursing, 22(1), 77-78.
3. Bellemare, S. (2007). Shaken baby syndrome vs inflicted brain injury. American Journal of Emergency Medicine, 25(9), 1080.
4. Best practices: Evidence-based nursing procedures (2nd ed.). (2006). Philadelphia, PA: Lippincott Williams and Wilkins.
5. Christian, C.W., Block, R., & the Committee on Child Abuse and Neglect. (2009). Abusive Head Trauma in Infants and Children. Pediatrics, 123(5), 1409-1411.
6. Fiske, E.A., & Hall, J.M. (2008). Inflicted childhood neurotrauma. ANS: Advances in Nursing Science, 31 (2), E1-8.
7. Mraz, M.A. (2009). The physical manifestations of shaken baby syndrome. Journal of Forensic Nursing, 5(1), 26-30.
8. Reynolds, A. (2008). Shaken baby syndrome: Diagnosis and treatment. Radiologic Technology, 80(2), 151-173.
9. Taketomo, C.K., Hodding, J.H., & Kraus, D. M. (2010). Pediatric dosage handbook (17th ed.). Hudson, OH: Lexi-Comp.



Department of «Pediatrics-1»

044-38

Methodological guidelines

7 page of 8



Department of «Pediatrics-1»

044-38

Methodological guidelines

8 page of 8