



ОҢТҮСТІК-ҚАЗАҚСТАН MEDISINA АКАДЕМИЯСЫ «Оңтүстік Қазақстан медицина академиясы» АҚ	 SKMA -1979-	SOUTH KAZAKHSTAN MEDICAL ACADEMY АО «Южно-Казахстанская медицинская академия»
Department of «Pediatrics-1» Methodological guidelines	044-38 1 page of 12	

**Methodological guidelines for teachers on teaching clinical skills in The center
of practical skills
(using action algorithms)**

Specialty: «General medicine» 6B10101
Discipline: «Basics of childhood diseases-1»
Course: 4th course
Department: Pediatrics-1
Compiler: Baltabaeva B.S.

Shymkent 2022

ОҢТҮСТІК-ҚАЗАҚСТАН MEDISINA AKADEMIASY «Оңтүстік Қазақстан медицина академиясы» АҚ	 SKMA -1979-	SOUTH KAZAKHSTAN MEDICAL ACADEMY АО «Южно-Казakhstanская медицинская академия»
Department of «Pediatrics-1» Methodological guidelines	044-38 2 page of 12	

Considered at a meeting of the Department of Pediatrics-1

Protocol №6 from 24.01.2023 year

Head of department, PhD  Kemelbekov K.S.

1. Name of clinical skill: Probe feeding of newborns.

2. Learning objective:

- To teach to correctly assess the degree of prematurity and severity of the condition to simulate a newborn in the gym;
- To teach students to work out practical skills on the technique of feeding newborns through the probe to automatism on simulators.
- Teach to choose the right way of feeding, which is determined by the severity of the child's condition, birth weight, gestational age;
- Use of enteral feeding in the maximum possible volume;
- Use only specialized formula milk for artificial feeding, intended for premature babies.

3. Pre-training and time to display the skill on the dummy: 10 min

4. Time to master the skill: 2 hours.

5. Theoretical knowledge required for mastering skills:


Indications. Unconsciousness of the child, mental disorders accompanied by complete refusal to eat, traumatic injuries of the oral cavity, lack of sucking or swallowing reflexes, prematurity of the III degree



Department of «Pediatrics-1»
Methodological guidelines

044-38
4 page of 12



ОҢТҮСТІК-ҚАЗАҚСТАН MEDISINA AKADEMIASY «Оңтүстік Қазақстан медицина академиясы» АҚ	 SKMA -1979-	SOUTH KAZAKHSTAN MEDICAL ACADEMY АО «Южно-Казахстанская медицинская академия»
Department of «Pediatrics-1» Methodological guidelines	044-38 5 page of 12	



Contraindications for probe feeding:

- malformations of the gastrointestinal tract, requiring emergency surgical intervention;
- abdominal distention;
- lack of active peristalsis;
- pathogenic meconium.

A gastric tube of appropriate size, which must meet the following requirements: be made of thermolabile material with special surface treatment for reduction of sticking effect in the mucosa to have a standard adapter of the type "Luer" for connection with a syringe or dropper, the cover on the adapter, colour code adaptor, centimeter scale, radiopaque strip. The recommended size of the probe depending on body mass

Recommended probe sizes depending on the child's body weight:

body weight less than-1000 g-nasal-No. 4, oral-No. 6, less than 2500 g-nasal-No. 6, oral-No. 10. Syringe (10 and 20 ml depending on the age and weight of the child).

Possible complications during the procedure of feeding through a gastric probe:


1. Using a probe of inappropriate size.

* Trauma of the nasal mucosa.

* Development of bedsores in the nasal passage.

* Hyperproduction of mucus with the development of secondary bacterial rhinitis.

2. "Tightening" of the probe in the oesophagus. At the same time, feeding by the free piston-free method is impossible, which is the main clinical criterion for the

ОҢТҮСТІК-ҚАЗАҚСТАН MEDISINA АКАДЕМИАСЫ «Оңтүстік Қазақстан медицина академиясы» АҚ	 SKMA -1979-	SOUTH KAZAKHSTAN MEDICAL ACADEMY АО «Южно-Казахстанская медицинская академия»
Department of «Pediatrics-1» Methodological guidelines	044-38 6 page of 12	

wrong position of the probe. However, if a dispenser or "piston" method is used, food will enter the esophagus, the following complications may develop:

- Injury of the esophagus
- Esophagitis
- * Vomiting with possible aspiration

In order to avoid such a complication, before each introduction of food, it is necessary to pull the piston on yourself and get the gastric contents.

3. Conducting a probe behind the pyloric flap into the duodenum. In this case, the food enters the small intestine, bypassing the stomach.

* Irritation of the pyloric sphincter, which leads to vomiting (the possibility of aspiration), a decrease or increase in the tone of the sphincter.

* Violation of the process of digestion, digestion and absorption.

6. A list of trainers, simulators, mannequins models:

1. Baby mannequin "Nursing Baby", without VitalSim^o
2. Mannequin baby CriSis, luxury with ECG (Deluxe Infant CriSis Manikin with EGG)
3. Mannequin baby CriSis, luxury with ECG (Deluxe Infant CriSis Manikin with EGG)

7. The list of medical products and equipment: simulator-1, gastric probe-1pc, glycerin - 1 PC, pink wipes 5-10, breast milk, phonendoscope-1pc, gloves-1 pair, adhesive plaster, 20 ml syringe, tray, centimeter tape.



8. Algorithm of execution skills

№	Algorithm of execution skills	Execution		
		Yes	No	Note
1	Wash and dry hands thoroughly, put on gloves.			
2	Measure the length of the insertion probe - from the earlobe to the tip of the nose and from the tip of the nose to the xiphoid process. To make a label on the probe			
3	To dial into the syringe the required amount of milk.			
4	Give the patient the correct position-lying on his side			
5	Take the probe with your right hand, pre-moistening the end of the probe in glycerin			
6	Insert the probe slowly through the nose or mouth to the mark.			
7	Make sure the correct introduction of the probe into the stomach (no choking, no wheezing during auscultation).			
8	Attach a syringe to the probe with the typed milk and make a slow introduction of milk			
9	Disconnect the syringe, close the end of the probe with a plug.			
10	Fix the probe with a band-aid on the child's cheek.			

9. Tasks

Task 1

The child was born from the fourth pregnancy, the second birth in the period of 27 weeks of gestation. Pregnancy proceeded against the background of toxicosis in the 2nd half, toxoplasmosis, fetal-placental insufficiency, anemia of medium degree. Body weight at birth 1050 grams, length 32 cm Diagnosed at birth: "RDS II degree." To 14 day life marks lethargy, muscular hypotension, oppression reflexes innate automatism, regurgitation, the absence of dynamics masses bodies, sucking reflex is missing. The skin is dry, icteric. In the lungs breathing weakened in all fields, no wheezing. There is a tendency to tachycardia, the rhythm is correct. The abdomen is soft, distended, peristalsis active.

Preliminary diagnosis: Bilateral pneumonia. Perinatal CNS lesion. Neonatal jaundice.


Your actions?

Demonstrate probe feeding on the simulator

Tests:

1. The length of the probe in the gastric method is:

A) 10-30 cm

ОҢТҰСТІК-ҚАЗАҚСТАН MEDISINA АКАДЕМИЯСЫ «Оңтүстік Қазақстан медицина академиясы» АҚ	 SKMA -1979-	SOUTH KAZAKHSTAN MEDICAL ACADEMY АО «Южно-Казахстанская медицинская академия»
Department of «Pediatrics-1» Methodological guidelines	044-38 8 page of 12	

B) 30-50 cm

C) 50-70 cm

D) 60-80 cm

E) 100 cm

2. The length of the probe in the duodenal method is:

A) 10-30 cm

C) 30-50 cm

C) 50-70 cm

D) 60-80 cm

E) 100 cm

3. The length of the probe inserted through the nose, a newborn with a body weight of 1000 g:

A) 13-21 cm

B) 21-26 cm

(C) 23-24 cm

D) 30-40 cm

E) 40-50 cm

4. The length of the probe inserted through the nose, a newborn with a body weight of 2000 g:

A) 13-21 cm

B) 21-26 cm

(C) 23-24 cm

D) 30-40 cm

E) 40-50 cm

5. The length of the probe, introduced through the nose, a newborn with a body weight of 2500-3500 g:

A) 13-21 cm

B) 21-26 cm

(C) 23-24 cm

D) 30-40 cm

E) 40-50 cm

6. The length of the probe inserted through the nose, a newborn with a body weight of 4000-5000 g:

A) 13-21 cm

B) 21-26 cm

(C) 23-24 cm

D) 30-40 cm

E) 40-50 cm

7. The inner diameter of the probe, administered to a newborn with a body weight of 1000 g:

- A) 0.64 mm
- C) 0.81 mm
- (C) 1 mm
- D) 2 mm
- E) 3 mm

8. The inner diameter of the probe administered to a newborn with a body weight of 2500-3500 g:

- A) 0.64 mm
- C) 0.81 mm
- (C) 1 mm
- D) 2 mm
- E) 3 mm


10. Evaluation criterions:

№	Execution	Evaluation criterions
1	Done	All steps are taken into account when performed correctly.
2	Half done	It is counted for small errors in the course of execution in 3 steps.
3	Not done	Makes mistakes during execution by more than 3 steps.

12. Literature:

Basic:

1. Geppe N. A. Propaedeutics of children's diseases. Moscow: GEOTAR-Media, 2012
2. Skvortsova V. A. Algorithms of feeding premature children: abstract. dis. ...Dr. med. sciences'. Moscow, 2002. 41 PP.
3. Skvortsova V. A., Borovik T. E., Yatsyk G. V. et al. Feeding premature children. 2006, No. 2, pp. 64-68.
4. Ntrebenko O. K. Some essential micronutrients in the nutrition of premature infants. Moscow, 2004. 136 PP.
5. Shabalov N. P. Neonatology. - In 2 vols. - M.: Medpress-inform, 2006.
6. Yatsyk G. V. Features of the digestive system in premature infants: abstract. dis. ... Dr. med. sciences'. M., 1980. 32 PP.

ОҢТҮСТІК-ҚАЗАҚСТАН MEDISINA АКАДЕМИЯСЫ «Оңтүстік Қазақстан медицина академиясы» АҚ	 SKMA -1979-	SOUTH KAZAKHSTAN MEDICAL ACADEMY АО «Южно-Казахстанская медицинская академия»
Department of «Pediatrics-1»	044-38	
Methodological guidelines	10 page of 12	

7. Sudareva O. A. Motor-evacuation function of the stomach in premature infants with different types of feeding: author. dis. ... Cand. honey. sciences'. M., 1985. 23 p.
8. Sadyrbaeva Z. S. Features of lipid metabolism in premature infants with different types of feeding: abstract. dis. ... Cand. honey. sciences'. M., 1982. 23 p.
9. Skvortsova V. A., Borovik T. E., Lukoyanova O. L., Gribakin S. G., Andreeva A.V. Modern tendencies of the problem of feeding premature children. *sovr. pediatrics*. 2005. Vol. 4, No. 2, pp. 80-84.

Additional:


1. Captain T. V. Propaedeutics of children's diseases with child care: textbook. 5th ed., extra.. - Moscow: Medpress-inform, 2009.
2. Berseth C. L. Effect of early feedings on maturation of the preterm infant's small intestine // *J. Pediatr*. 1992. Vol. 120, pp. 947-953.
3. Dusik A. M., Poindexter, B. B., Ehenkranz R. A. et al. Growth failure in the preterm infant: can we catch up? *Semin.Perinatol.*, 2003, 27 (4), pp. 302-310.
4. Micheli J. L., Fawer C. L., Schutz Y. Protein some requirement of the extremely low birthweight preterm infant // 43-rd Nestle Nutrition Workshop. *Nutrition of the Very Low Birth Weight Infant*. - Warsaw, 1998, 30 s.
5. Lucas A. *Clinical Nutrition of the Young Child*. Feeding the preterm infants. New York, 1991, p. 311-355.
6. Koletzko B. Long-chain polyunsaturated fatty acids in the diets of premature infants / *Polyunsaturated fatty acids in human nutrition*. Nestle Nutrition Workshop Series. 1992 Vol.28, p. 135-142.
7. Owners L., Burrin D., Berseth C. L. Enteral nutrition has a dose-response effect on maturation of neonatal canine motor activity / *Gastroenterology*. 1996. Vol. 110, p. 828.
8. Lucas A. Nutrition, growth and development of postdischarge preterm infants. In: *Posthospital nutrition in the preterm infant: Report of the 106th Ross Conference on Pediatric Research*. 1996, p. 81-89.
9. Lucas A., Fewtrell M. S., Morley R. et. al. Randomized trial of nutrient-enriched formula versus standard formula for post discharge preterm infants. *Pediatr*. 2001; 108 (3), p. 703-711.


Electronic resources:

1. Lissauer, T. *Childhood illnesses [Electronic resource] : textbook* / T. Lissauer, G. Clayden ; lane. from English. N. A. Geppe. - Electron. text messages. 329mb). - Moscow: GEOTAR-Media, 2017. - el. wholesale. disc (CD-ROM).

13. Standards of correct answers for the assessment material:

Answers: 1-B, 2-C, 3-A, 4-B, 5-C, 6-D, 7-B, 8-C.

<p>ОҢТҮСТІК-ҚАЗАҚСТАН MEDISINA AKADEMIASY «Оңтүстік Қазақстан медицина академиясы» АҚ</p>		<p>SOUTH KAZAKHSTAN MEDICAL ACADEMY АО «Южно-Казakhstanская медицинская академия»</p>
<p>Department of «Pediatrics-1»</p>		<p>044-38</p>
<p>Methodological guidelines</p>		<p>11 page of 12</p>

<p>ОҢТҮСТІК-ҚАЗАҚСТАН MEDISINA AKADEMIASY «Оңтүстік Қазақстан медицина академиясы» АҚ</p>		<p>SOUTH KAZAKHSTAN MEDICAL ACADEMY АО «Южно-Казakhstanская медицинская академия»</p>
<p>Department of «Pediatrics-1»</p>		<p>044-38</p>
<p>Methodological guidelines</p>		<p>12 page of 12</p>